

ZFIRE: SIMILAR STELLAR GROWTH IN H α -EMITTING CLUSTER AND FIELD GALAXIES AT $z \sim 2$

KIM-VY H. TRAN^{1,2}, LEO Y. ALCORN², GLENN G. KACPRZAK³, THEMIYA NANAYAKKARA³, CAROLINE STRAATMAN⁴,
TIAN TIAN YUAN⁵, MICHAEL COWLEY^{6,7}, ROMÉEL DAVÉ^{8,9,10}, KARL GLAZEBROOK³, LISA J. KEWLEY⁵, IVO LABBÉ⁴, DAVIDÉ
MARTIZZI¹¹, CASEY PAPOVICH², RYAN QUADRI², LEE R. SPITLER^{6,7}, ADAM TOMCZAK¹²

¹kimvy.tran@tamu.edu

²George P. and Cynthia W. Mitchell Institute for Fundamental Physics and Astronomy, Department of Physics & Astronomy, Texas A&M University, College Station, TX 77843

³Swinburne University of Technology, Hawthorn, VIC 3122, Australia

⁴Leiden Observatory, Leiden University, P.O. Box 9513, NL-2300 RA Leiden, The Netherlands

⁵Research School of Astronomy and Astrophysics, The Australian National University, Cotter Road, Weston Creek, ACT 2611, Australia

⁶Department of Physics and Astronomy, Faculty of Science and Engineering, Macquarie University, Sydney, NSW 2109, Australia

⁷Australian Astronomical Observatory, PO Box 915, North Ryde, NSW 1670, Australia

⁸University of the Western Cape, Bellville, Cape Town, 7535, South Africa

⁹South African Astronomical Observatories, Observatory, Cape Town, 7925, South Africa

¹⁰African Institute for Mathematical Sciences, Muizenberg, Cape Town, 7945, South Africa

¹¹Department of Astronomy, University of California, Berkeley, CA, 94720

¹²Department of Physics, University of California, Davis, CA, 95616

ABSTRACT

We compare galaxy scaling relations as a function of environment at $z \sim 2$ with our ZFIRE survey^a where we have measured H α fluxes for 90 star-forming galaxies selected from a mass-limited [$\log(M_*/M_\odot) > 9$] sample based on ZFOURGE^b. The cluster galaxies (37) are part of a confirmed system at $z = 2.095$ and the field galaxies (53) are at $1.9 < z < 2.4$; all are in the COSMOS legacy field. There is no statistical difference between H α -emitting cluster and field populations when comparing their star formation rate (SFR), stellar mass (M_*), galaxy size (r_{eff}), SFR surface density [$\Sigma(\text{H}\alpha_{\text{star}})$], and stellar age distributions. The only difference is that at *fixed stellar mass*, the H α -emitting cluster galaxies are $\log(r_{\text{eff}}) \sim 0.1$ larger than in the field. Approximately 19% of the H α -emitters in the cluster and 26% in the field are IR-luminous ($L_{\text{IR}} > 2 \times 10^{11} L_\odot$). Because the LIRGs in our combined sample are ~ 5 times more massive than the low-IR galaxies, their radii are $\sim 70\%$ larger. To track stellar growth, we separate galaxies into those that lie above, on, and below the H α star-forming main sequence (SFMS) using $\Delta\text{SFR}(M_*) = \pm 0.2$ dex. Galaxies above the SFMS (starbursts) tend to have higher H α SFR surface densities and younger light-weighted stellar ages compared to galaxies below the SFMS. Our results indicate that starbursts (+SFMS) in the cluster and field at $z \sim 2$ are growing their stellar cores. Lastly, we compare to the (SFR– M_*) relation from RHAPSODY-G cluster simulations and find the predicted slope is nominally consistent with the observations. However, the predicted cluster SFRs tend to be too low by a factor of ~ 2 which seems to be a common problem for simulations across environment.

Keywords: galaxies: evolution – galaxies: star formation – galaxies: starburst – galaxies: structure – galaxies: clusters: individual (COSMOS) – infrared: galaxies

1. INTRODUCTION

With the discovery and spectroscopic confirmation of galaxy clusters at $z \sim 2$, we have reached the epoch when many massive galaxies in clusters are still forming a significant fraction of their stars (*e.g.* Tran et al. 2010; Papovich et al. 2010; Zeimann et al. 2012; Brodwin et al. 2013; Gobat et al. 2013; Webb et al. 2015). We can now pinpoint when cluster galaxies begin to diverge from their field counterparts and thus separate evolution driven by galaxy mass

^a <http://zfire.swinburne.edu.au>

^b <http://zfourge.tamu.edu>

from that of environment (Peng et al. 2010; Wetzel et al. 2012; Quadri et al. 2012; Muzzin et al. 2012; Papovich et al. 2012; Bassett et al. 2013). At this epoch, measurements of galaxy properties such as stellar mass, star formation rate, physical size, and metallicity have added leverage because the cosmic star formation rate density peaks at $z \sim 2$ (see review by Madau & Dickinson 2014, and references therein). Observed galaxy scaling relations also test current formation models (e.g. Davé et al. 2011; Genel et al. 2014; Tonnesen & Cen 2014; Schaye et al. 2015; Hahn et al. 2015; Martizzi et al. 2016).

Particularly useful for measuring galaxy scaling relations at $z \sim 2$ are mass-limited surveys because they bridge UV/optical selected galaxies with the increasing number at $z \gtrsim 2$ of dusty star-forming systems that are IR-luminous but UV faint (see reviews by Lutz 2014; Casey et al. 2014, and references therein). Large imaging surveys have measured sizes and morphologies for galaxies (e.g. Wuyts et al. 2011; van der Wel et al. 2012), but these studies use photometric redshifts based on broad-band photometry and are limited to $\log(M_*/M_\odot) \gtrsim 10$ at $z \sim 2$, i.e. just below the characteristic stellar mass at this epoch (Tomczak et al. 2014). Pushing to lower stellar masses at $z \sim 2$ with more precise star formation rates requires deep imaging that spans rest-frame UV to near-IR wavelengths to fully characterize the galaxy Spectral Energy Distributions (SEDs) and obtain reliable photometric redshifts and stellar masses (Brammer et al. 2008, 2012; Brown et al. 2014; Forrest et al. 2016).

Here we combine H α emission from our ZFIRE survey (Nanayakkara et al. 2016) with galaxy properties from the ZFOURGE survey (Straatman et al. 2016) with IR luminosities from *Spitzer* to track how galaxies grow at $z \sim 2$. ZFIRE is a near-IR spectroscopic survey with MOSFIRE (McLean et al. 2012) on Keck I where targets are selected from ZFOURGE, an imaging survey that combines deep near-IR observations taken with the FourStar Imager (Persson et al. 2013) at the *Magellan Observatory* with public multi-wavelength observations, e.g. *Hubble Space Telescope* imaging from CANDELS (Grogin et al. 2011). Because ZFIRE is based on ZFOURGE which is mass-complete to $\log(M_*/M_\odot) \sim 9$ at $z \sim 2$ (Tomczak et al. 2014; Straatman et al. 2016), we can measure galaxy scaling relations for cluster and field galaxies spanning a wide range in stellar mass.

With spectroscopic redshifts and deep multi-wavelength coverage, we also are able to compare IR luminous to low-IR galaxies in one of the deepest mass-limited studies to date. Swinbank et al. (2010) find that submillimeter galaxies (among the dustiest star-forming systems in the universe) at $z \sim 2$ have similar radii in the rest-frame optical as “normal” star-forming field galaxies, but Kartaltepe et al. (2012) find that Ultra Luminous Infra-Red Galaxies (ULIRGs; $L_{\text{IR}} > 10^{12} L_\odot$) at $z \sim 2$ have larger radii than typical galaxies. In contrast, Rujopakarn et al. (2011) find that local ULIRGs have smaller radii than the star-forming field galaxies. Because of these conflicting results, it is still not clear whether the IR-luminous phase for star-forming galaxies at $z \sim 2$ is correlated with size growth.

Alternatively, a more effective approach may be to consider galaxies in terms of their star formation rate versus stellar mass, i.e. the Star-Forming Main Sequence (SFMS; Noeske et al. 2007; Whitaker et al. 2014; Tomczak et al. 2016, and numerous other studies). For example, Wuyts et al. (2011) find that galaxies above the SFMS tend to have smaller effective radii. By separating galaxies into those above, on, and below the SFMS, recent studies find that galaxy properties such as Sérsic index and gas content correlate with a galaxy’s location relative to the SFMS (Genzel et al. 2015; Whitaker et al. 2015). However, these studies use star formation rates based on SED fits to rest-frame UV-IR observations. Here we explore these relations using H α to measure the instantaneous SFRs of galaxies at $z \sim 2$.

We focus on the COSMOS legacy field where we have identified and spectroscopically confirmed a galaxy cluster at $z = 2.095$ (hereafter the COSMOS cluster; Spitler et al. 2012; Yuan et al. 2014). We build on our ZFIRE results comparing the cluster to the field for the gas phase metallicity- M_* relation (Kacprzak et al. 2015, 2016), the ionization properties of the Inter-Stellar Medium (ISM; Kewley et al. 2016), and the kinematics and virial masses of individual galaxies (Alcorn et al. 2016). There are also a number of luminous infrared sources that are likely dusty star-forming galaxies in the larger region around the COSMOS cluster (Hung et al. 2016).

We use a Chabrier Initial Mass Function and AB magnitudes throughout our analysis. We assume $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$, and $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$. At $z = 2$, the angular scale is $1'' = 8.37 \text{ kpc}$.

2. OBSERVATIONS

2.1. ZFOURGE Catalog

To select spectroscopic targets in the COSMOS field, we use the ZFOURGE catalog that provides high accuracy photometric redshifts based on multi-filter ground and space-based imaging (Straatman et al. 2016). ZFOURGE uses EAZY (Brammer et al. 2008, 2012) to first determine photometric redshifts by fitting Spectral Energy Distributions, and then FAST (Kriek et al. 2009) to measure rest-frame colors, stellar masses, stellar attenuation, and specific star formation rates for a given SF history. We use a Chabrier (2003) initial stellar mass function, constant solar metallicity,

and exponentially declining star formation rate ($\tau = 10$ Myr to 10 Gyr). For a detailed description of the ZFOURGE survey and catalogs, we refer the reader to [Straatman et al. \(2016\)](#).

An advantage of using the deep ZFOURGE catalog is that we can optimize the target selection to MOSFIRE, specifically by selecting star-forming galaxies as identified by their UVJ colors (*e.g.* [Wuyts et al. 2007](#); [Williams et al. 2009](#)). Because the ZFOURGE catalog reaches Fourstar/Ks = 25.3 mag and fits the SEDs from the UV to MIR ([Straatman et al. 2016](#)), we are able to obtain MOSFIRE spectroscopy for objects with stellar masses down to $\log(M_*/M_\odot) \sim 9$ at $z \sim 2$ ([Nanayakkara et al. 2016](#)). Our analysis focuses on the star-forming galaxies, thus we remove AGN identified in [Cowley et al. \(2016\)](#)’s multi-wavelength catalog.

2.2. Keck/MOSFIRE Spectroscopy

We refer the reader to [Nanayakkara et al. \(2016\)](#) and [Tran et al. \(2015\)](#) for an extensive description of our Keck/MOSFIRE data reduction and analysis. To briefly summarize, the spectroscopy was obtained on observing runs in December 2013 and February 2014. A total of eight slit-masks were observed in the K-band with total integration time of 2 hours each. The K-band wavelength range is $1.93 - 2.38 \mu\text{m}$ and the spectral dispersion of $2.17 \text{ \AA/pixel}$. We also observed two masks in H-band covering $1.46 - 1.81 \mu\text{m}$ with a spectral dispersion $1.63 \text{ \AA/pixel}$.

To reduce the MOSFIRE spectroscopy, we use the publicly available data reduction pipeline (DRP) developed by the instrument team¹. We then apply custom IDL routines to correct the reduced 2D spectra for telluric absorption, spectro-photometrically calibrate by anchoring to the well-calibrated photometry, and extract the 1D spectra with associated 1σ error spectra (see [Nanayakkara et al. 2016](#)). We reach a line-flux of $\sim 0.3 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$ (5σ ; [Nanayakkara et al. 2016](#)). In our analysis, we select galaxies with H α redshifts of $1.9 < z < 2.4$, *i.e.* corresponding to the K-band wavelength range, and exclude AGN (3 in cluster, 6 in field) identified by [Cowley et al. \(2016\)](#).

As reported in [Nanayakkara et al. \(2016\)](#), our success rate of detecting H α emission at $S/N > 5$ in the K-band is $\sim 73\%$ and the redshift distribution of the H α -detected galaxies is the same as the expected redshift probability distribution from ZFOURGE (see their Fig. 6). A higher success rate is nearly impossible given the number of strong sky-lines within the K-band. We also confirm that the ZFIRE galaxies are not biased in stellar mass compared to the ZFOURGE photometric sample ([Nanayakkara et al. 2016](#), see their §3.3 & Fig. 8).

Figure 1 shows the spatial distribution of our 37 cluster and 53 field galaxies at $z \sim 2$. Cluster members have spectroscopic redshifts of $2.08 < z_{\text{spec}} < 2.12$ ([Yuan et al. 2014](#); [Nanayakkara et al. 2016](#)) and field galaxies have z_{spec} of $(1.97 - 2.06)$ and $(2.13 - 2.31)$. We consider only galaxies with z_{spec} quality flag $Q_z = 3$. To test if our field sample is contaminated by cluster galaxies, we also apply a more stringent redshift selection of $(1.97 - 2.03)$ and $(2.17 - 2.31)$ which corresponds to > 8 times the cluster’s velocity dispersion from the cluster redshift ($\sigma_{1D} = 552 \text{ km s}^{-1}$; [Yuan et al. 2014](#)). We confirm that using the more conservative redshift range for the field does not change our following results.

We note that our study focuses on cluster and field galaxies at $z \sim 2$ identified by their H α emission, thus we cannot confidently measure the relative fraction of star-forming to all galaxies across environment with the current dataset.

2.3. Measuring Galaxy Sizes & Morphologies

We use GALFIT ([Peng et al. 2010](#)) to measure Sérsic indices, effective radii, axis ratios, and position angles for the spectroscopically confirmed galaxies in COSMOS using *Hubble Space Telescope* imaging taken with WFC3/F160W. Most of these galaxies are in the [van der Wel et al. \(2012\)](#) morphological catalog which spans a wide redshift range. However, we choose to measure independently the galaxy sizes and morphologies to optimize the fits for our galaxies at $z \sim 2$.

Of the 90 galaxies in our H α -emitting sample, we measure effective radii along the major axis and Sérsic indices for 83 (35 cluster, 48 field); see Figs. 2 & 3 for galaxy images and Table 1 for galaxy properties. Seven of the galaxies could not be fit because of contamination due to diffraction spikes from nearby stars or incomplete F160W imaging (see [Skelton et al. 2014](#)). We include a quality flag on the GALFIT results and identify 12 galaxies with fits that have large residuals due to, *e.g.* being mergers (see [Alcorn et al. 2016](#)). We confirm that excluding these 12 galaxies does not change our general results and so we use the effective radii measured for all 83 galaxies in our analysis.

Following [van der Wel et al. \(2014\)](#), we use the effective radius to characterize size because r_{eff} is more appropriate than a circularized radius for galaxies spanning the range in axis ratios. We confirm that using r_{circ} instead of r_{eff} does not change the following results except for shifting the size distribution of the entire galaxy sample to smaller sizes.

¹ <https://github.com/Mosfire-DataReductionPipeline/MosfireDRP>




Figure 1. Spatial distribution of $H\alpha$ -emitting cluster (filled circles; 37) and field (crosses; 53) galaxies at $z \sim 2$ in the COSMOS legacy field. Galaxies with total IR luminosities $L_{\text{IR}} > 2 \times 10^{11} L_{\odot}$ as measured using *Spitzer*/ $24\mu\text{m}$ (3σ detection) are shown as open stars (21). Active Galactic Nuclei (AGN) are excluded using the AGN catalog by [Cowley et al. \(2016\)](#). The fraction of IR-luminous galaxies is the same in the field and the cluster ($\sim 20 - 25\%$).

The trends in the scaling relations that depend on galaxy size, *e.g.* comparing cluster to field and galaxies relative to the SFMS, are robust.

2.4. Dust-corrected $H\alpha$ Star Formation Rates

To use $H\alpha$ line emission as a measure of star formation rate (SFR), we need to correct for dust attenuation. Although determining the internal extinction using the Balmer decrement is preferred, we have $H\beta$ for only a small subset. Thus we must rely on the stellar attenuation $A_{V,\text{star}}$ measured by FAST which assumes $R_V=4.05$ (starburst attenuation curve; [Calzetti et al. 2000](#))². For more extensive results on stellar vs. Balmer-derived attenuation and star formation rates, we refer the reader to [Price et al. \(2014\)](#) and [Reddy et al. \(2015\)](#).

Following [Tran et al. \(2015\)](#) (see also [Steidel et al. 2014](#)), the $H\alpha$ line fluxes are corrected using the nebular attenuation curve from [Cardelli et al. \(1989\)](#) with $R_V=3.1$

² The starburst (SB) attenuation curve is commonly referred to as the Calzetti law and is appropriate for continuum measurements. We use “starburst” as requested by D. Calzetti.




Figure 2. HST images ($4'' \times 4''$) generated by summing F125W, F140W, and F160W for $\text{H}\alpha$ -emitting cluster galaxies ($2.08 < z_{\text{spec}} < 2.12$); Sérsic indices and effective radii are measured using GALFIT for 35 of 37 members. Galaxies are labeled with their ZFIRE IDs and IR-luminous galaxies ($L_{\text{IR}} > 2 \times 10^{11} L_{\odot}$) are noted as LIRGs.




Figure 3. HST images ($4'' \times 4''$) generated by summing F125W, F140W, and F160W for $\text{H}\alpha$ -emitting field galaxies at $z \sim 2$ ($1.9 < z_{\text{spec}} < 2.4$); Sérsic indices and effective radii are measured using GALFIT for 49 of 53 field galaxies. Galaxies are labeled with their ZFIRE IDs and IR-luminous galaxies ($L_{\text{IR}} > 2 \times 10^{11} L_{\odot}$) are noted as LIRGs.

$$A(\text{H}\alpha)_{\text{HII}} = 2.53 \times E(\text{B} - \text{V})_{\text{HII}}. \quad (1)$$

We use the observed stellar to nebular attenuation ratio of $E(\text{B} - \text{V})_{\text{star}} = 0.44 \times E(\text{B} - \text{V})_{\text{HII}}$ (Calzetti et al. 2000) and the color excess $E(\text{B} - \text{V})_{\text{star}}$ which is the stellar attenuation $A_{\text{V,star}}$ measured by FAST divided by $R_{\text{V}} = 4.05$. Combining these factors, we have

$$A(\text{H}\alpha)_{\text{HII}} = 5.75 \times E(\text{B} - \text{V})_{\text{star}} \quad (2)$$

which we use to correct all of the $\text{H}\alpha$ fluxes for attenuation. Recent work by Reddy et al. (2015) suggests that the $E(\text{B} - \text{V})_{\text{star}}$ to $E(\text{B} - \text{V})_{\text{HII}}$ ratio may depend on stellar mass at $z \sim 2$, but there is significant scatter in the fitted

relation. We stress that such a correction would not change our results because we use the same method to measure $H\alpha$ -SFRs for all the galaxies in our study and compare internally.

We determine the corresponding star formation rates using the relation from Hao et al. (2011):

$$\log[\text{SFR}(H\alpha_{\text{star}})] = \log[L(H\alpha_{\text{star}})] - 41.27 \quad (3)$$

This relation assumes a Kroupa IMF ($0.1 - 100 M_{\odot}$; Kroupa 2001), but the relation for a Chabrier IMF is virtually identical (difference of 0.05). Note that $\log[\text{SFR}(H\alpha_{\text{star}})]$ values determined with the Hao et al. (2011) relation are 0.17 dex lower than when using the Kennicutt (1998) relation.

2.5. $H\alpha_{\text{star}}$ -SFR Surface Densities

With the $H\alpha_{\text{star}}$ SFRs and galaxy sizes as measured by their effective radii (r_{eff}), we can then determine the SFR surface density:

$$\Sigma(H\alpha_{\text{star}}) = \frac{\text{SFR}(H\alpha_{\text{star}})}{2\pi \times r_{\text{eff}}^2} \quad (4)$$

Note that most of the cluster and field galaxies have effective radii of $r_{\text{eff}} \sim 0.35''$ (Fig. 4) which is comparable to the slit-width of $0.7''$.

It is possible that by using r_{eff} measured with WFC/F160W imaging, we are overestimating $\Sigma(H\alpha_{\text{star}})$. Förster Schreiber et al. (2011) find that the $H\alpha$ sizes of six $z \sim 2$ galaxies are comparable to their rest-frame continuum sizes as measured with IFU and HST observations. However, Nelson et al. (2016) show that at $z \sim 1$, continuum-based sizes tend to be smaller than $H\alpha$ -based sizes for star-forming galaxies with $\log(M_{\star}/M_{\odot}) \gtrsim 10$. While correcting for a possible dependence of $H\alpha$ -size on galaxy mass would shift $\Sigma(H\alpha_{\text{star}})$ to lower values, it would not change our overall conclusions based on comparing the different galaxy populations.

Note that with our current single-slit observations, we cannot address a possible environmental dependence of $H\alpha$ -disks. Galaxies in the Virgo cluster are known to have truncated $H\alpha$ -disks compared to the field (Kenney & Koopmann 1999; Koopmann & Kenney 2004), thus not accounting for disk truncation in the cluster galaxies may lead to overestimating their total $H\alpha_{\text{star}}$ -SFRs and consequently $\Sigma(H\alpha_{\text{star}})$. Future deep IFU observations with the next generation of large telescopes should be able to test for $H\alpha$ -disk truncation in these $z \sim 2$ galaxies.

2.6. IR Luminosities from Spitzer/MIPS

Summarizing from Tomczak et al. (2016), IR luminosities are determined from Spitzer/MIPS observations at $24\mu\text{m}$ (GOODS-S: PI M. Dickinson, COSMOS: PI N. Scoville, UDS: PI J. Dunlop) which have 1σ uncertainties of $10.3 \mu\text{Jy}$ in COSMOS. We measure the $24\mu\text{m}$ fluxes within $3.5''$ apertures and use the custom code MOPHONGO (written by I. Labbé; see Labbé et al. 2006; Wuyts et al. 2007) to deblend fluxes from multiple sources. The Wuyts et al. (2008) templates are fit to the SEDs using the $H\alpha$ -redshifts to determine integrated $8 - 1000 \mu\text{m}$ fluxes; we refer the reader to Tomczak et al. (2016) for a full description of the IR measurements.

For galaxies at $z \sim 2$, the 3σ L_{IR} detection limit is $2 \times 10^{11} L_{\odot}$, *i.e.* all our L_{IR} galaxies are Luminous Infra-Red Galaxies³. Figure 1 shows the spatial distribution of IR-luminous cluster and field galaxies. In our analysis, we use IR-based luminosities and $H\alpha_{\text{star}}$ SFRs. We note that L_{IR} detection thresholds at $z > 1$ correspond to star formation rates that are much higher than UV-based SFRs. Thus comparing, *e.g.* $H\alpha_{\text{star}}$ -SFR to a combined (IR+UV) SFR instead of an L_{IR} -only SFR does not change our results.

3. RESULTS

3.1. A Population of IR-Luminous Galaxies

A remarkable 19% (7/37) of $H\alpha$ -emitting cluster galaxies at $z \sim 2$ have $L_{\text{IR}} > 2 \times 10^{11} L_{\odot}$. Within errors, this fraction of IR-luminous cluster galaxies is comparable to the field (26%, 14/53; Fig. 1). Saintonge et al. (2008) showed using $24\mu\text{m}$ observations of ~ 1500 spectroscopically confirmed cluster galaxies that the fraction of IR members increases with redshift, but this was limited to galaxy clusters at $0 < z < 1$. More recent studies using the *Herschel Space Observatory* have detected IR sources in galaxy clusters at $z > 1$ (Popesso et al. 2012; Santos et al. 2014), but far-IR observations can only detect a handful of the most IR-luminous systems with star formation rates $> 100 M_{\odot} \text{ yr}^{-1}$.

³ Note that our L_{IR} detection limit is higher than the LIRG threshold of $10^{11} L_{\odot}$ (see review by Sanders & Mirabel 1996), *i.e.* we do not detect LIRGs with ($10^{11} L_{\odot} < L_{\text{IR}} < 2 \times 10^{11} L_{\odot}$). Thus some of our low-IR galaxies may still technically be LIRGs




Figure 4. We measure the effective radii (r_{eff}) using *Hubble Space Telescope* imaging taken with WFC3/F160W. *Left:* The galaxy size-stellar mass relation for our combined sample is consistent with the fit to star-forming galaxies at $z \sim 2$ measured using photometric redshifts by CANDELS and clearly offset from the relation at $z = 0.25$ (pink dash-dot curves; van der Wel et al. 2014). We find no significant difference between the size-mass relation for H α -emitting cluster (red dashed) and field (blue dotted) galaxies at $z \sim 2$. *Right:* The r_{eff} - M_* relations for galaxies on (open crosses) and below (filled triangles) the H α star-forming main sequence (SFMS; see Fig. 6) are consistent with CANDELS, but the galaxies with elevated SFRs (filled squares) have smaller radii at a given stellar mass. For reference, the black line is the (2σ clipped) least squares fit to our combined sample.

Our survey is the first to spectroscopically confirm the high fraction of LIRGs in galaxy clusters at $z \sim 2$ (see also Hung et al. 2016).

3.2. Comparing Star Formation Rates

3.2.1. Cluster vs. Field

We find no evidence of different correlations between H α and L_{IR} when considering the cluster and field samples separately (Fig. 5; Table 1). For the 14 field and 7 cluster galaxies with $L_{\text{IR}} > 2 \times 10^{11} L_{\odot}$, a K-S test measures a p-value of 0.13, *i.e.* the statistical likelihood of the cluster and field populations being drawn from different parent populations is low. The average $\log(L_{\text{IR}})$ per galaxy is comparable: 11.7 ± 0.3 in the field versus 11.8 ± 0.3 in the cluster. This is true also when selecting instead by $\text{SFR}(\text{H}\alpha_{\text{star}}) > 2 M_{\odot} \text{ yr}^{-1}$: the field (52) and cluster (34) populations have the same median $\log[\text{SFR}(\text{H}\alpha_{\text{star}})]$ of 0.9 ± 0.3 . Note that K-S tests confirm the H α -emitting galaxies in the cluster and field are drawn from the same parent population in terms of their stellar mass and Specific Star Formation Rate ($\text{SSFR} = \text{SFR}/M_*$).

3.2.2. H α vs. L_{IR}

For galaxies with both $\text{H}\alpha_{\text{star}} > 2 M_{\odot} \text{ yr}^{-1}$ and $L_{\text{IR}} > 2 \times 10^{11} L_{\odot}$ (21), a Spearman rank test confirms a positive correlation ($> 2\sigma$) between SFRs based on these two tracers (Fig. 5, Table 1; see also Ibar et al. 2013; Shivaee et al. 2016). However, the dust-corrected $\text{H}\alpha_{\text{star}}$ SFRs are systematically lower than L_{IR} SFRs by ~ 0.5 dex, *i.e.* by nearly a factor of 3. This is driven mostly by a combination of using the Hao et al. (2011) relation for converting H α luminosities to SFRs instead of, *e.g.* Kennicutt (1998), and by choice of dust law. We confirm that comparing $\text{H}\alpha_{\text{star}}$ to a combined (IR+UV) star formation rate does not change our results.

We measure a scatter of $\sigma \sim 0.33$ dex in $\text{H}\alpha_{\text{star}}$ - L_{IR} SFRs which is larger than $\sigma \sim 0.22$ dex measured recently by Shivaee et al. (2016) for 17 galaxies at $z \sim 2$. However, their analysis focuses on galaxies with SFRs $> 10 M_{\odot} \text{ yr}^{-1}$ while we push to $\text{H}\alpha_{\text{star}}$ SFRs of $\sim 2 M_{\odot} \text{ yr}^{-1}$. From Fig. 5, the discrepancy between $\text{H}\alpha_{\text{star}}$ and L_{IR} SFRs decreases at higher values.

3.3. H α Star-Forming Main Sequence at $z \sim 2$




Figure 5. A Spearman rank test confirms that for the 21 galaxies with $\text{SFR}(\text{H}\alpha_{\text{star}}) > 2 \text{ M}_{\odot} \text{ yr}^{-1}$ and $L_{\text{IR}} > 2 \times 10^{11} L_{\odot}$ (horizontal dotted line), their SFRs based on these two tracers are correlated ($> 2\sigma$ confidence). The solid line shows the best fit least squares (2σ -clipped) and the dashed diagonal line is parity; the cross in the upper left shows a representative log error of ± 0.1 dex. Galaxies with $L_{\text{IR}} < 2 \times 10^{11} L_{\odot}$ are shown in gray and have L_{IR} errors larger than the representative value. There is no evidence of environmental dependence: K-S tests confirm that the $\text{H}\alpha_{\text{star}}$ and L_{IR} star formation rates have the same parent populations for cluster and field galaxies. The same is true if we compare the combined (IR+UV) star formation rate to $\text{H}\alpha_{\text{star}}$ values. However, SFRs based on $\text{H}\alpha_{\text{star}}$ are systematically lower than those from L_{IR} .

Using deep multi-wavelength imaging, the relation between star formation rate and stellar mass is now measured to $z \sim 3$ for thousands of galaxies down to $\log(M_{\star}/M_{\odot}) \sim 9$ (e.g. Whitaker et al. 2012; Tomczak et al. 2016, see Fig. 6). However, the SFRs and stellar masses derived by fitting Spectral Energy Distributions (SEDs) to multi-wavelength imaging can be degenerate. Measurements of $\text{H}\alpha$ fluxes are a more accurate tracer of the instantaneous SFR than fitting SEDs to photometry (Kennicutt & Evans 2012), but are restricted to a smaller sample of galaxies due to the observational challenge of measuring $\text{H}\alpha$ at $z \sim 2$.

Combining SFRs based on $\text{H}\alpha_{\text{star}}$ fluxes and stellar masses derived from SED fitting, we fit the $\text{SFR}-M_{\star}$ relation using a (2σ -clipped) least-squares for the field and cluster populations separately. Note that the field and cluster galaxies span the full range in both stellar mass and $\text{H}\alpha_{\text{star}}$ -SFR (Fig. 6). The cluster and field galaxies at $z \sim 2$ have the same increasing $\text{SFR}-M_{\star}$ relation:

$$\log[\text{SFR}(\text{H}\alpha_{\text{star}}, \text{Field})] = 0.69 \log(M_{\star}) - 5.82 \quad (5)$$




Figure 6. *Left:* At $z \sim 2$, the COSMOS cluster (red filled circles) and field (blue line stars) galaxies follow identical relations between stellar mass and $H\alpha_{\text{star}}$ star formation rate; 2σ -clipped least-squares fits are shown by red dashed and blue dotted lines, respectively. The cross in the lower right shows a representative log error of ± 0.1 dex. Both fits are consistent with the shape of the SFR- M_* relation measured by ZFOURGE for star-forming field galaxies at $z \sim 2$ using photometric redshifts (pink curve Tomczak et al. 2016) as well as the mass-binned sample from MOSDEF for $H\alpha$ -selected field galaxies at $z \sim 2$ (open triangles; Sanders et al. 2015); because we use Hao et al. (2011) to convert $H\alpha$ luminosity to SFR, we are offset in $\log[\text{SFR}(H\alpha_{\text{star}})]$ from both ZFOURGE and MOSDEF. The more massive galaxies [$\log(M_*/M_\odot) > 10$] tend to be IR-luminous ($L_{\text{IR}} > 2 \times 10^{11} L_\odot$), *i.e.* they are LIRGs. *Right:* We fit the $H\alpha$ star-forming main sequence (hereafter SFMS) using our combined cluster and field sample (cyan line). In our analysis, we consider star-forming galaxies that lie above (+SFMS; purple filled squares), on (=SFMS; cyan open crosses), and below (-SFMS; yellow filled triangles) the $H\alpha$ SFMS. Also shown is the predicted SFMS relation at $z \sim 2$ from RHAPSODY-G, a high-resolution AMR simulation of galaxy clusters (gray long dash-dot line; Martizzi et al. 2016).

$$\log[\text{SFR}(H\alpha_{\text{star}}, \text{Cluster})] = 0.62 \log(M_*) - 5.15 \quad (6)$$

$$\log[\text{SFR}(H\alpha_{\text{star}}, \text{All})] = 0.61 \log(M_*) - 5.11 \quad (7)$$

where SFR is in $M_\odot \text{ yr}^{-1}$ and M_* is in $M_\odot$. The rms error on the fitted slopes is ~ 0.2 , and separate 1D K-S tests confirm that the stellar mass and SFR distributions of our cluster and field populations are similar. A possible concern is that our field sample could be contaminated by cluster members, but we confirm that applying a more stringent redshift cut of $> 8\sigma_{1D}$ to select field galaxies does not change our results.

Our measurements are consistent with recent results, *e.g.* from ZFOURGE (SED fitting of UV-MIR; Tomczak et al. 2016) and MOSDEF ($H\alpha$; Sanders et al. 2015), and span similar ranges in stellar mass and star formation rate. However, our $H\alpha_{\text{star}}$ SFRs are lower. This offset is mostly likely due to differences in the relation used to convert $H\alpha$ luminosities to SFRs, *e.g.* Hao et al. (2011) vs. Kennicutt (1998), and choice of dust law. Accounting for both these effects increases $\log[\text{SFR}(H\alpha_{\text{star}})]$ by ~ 0.3 dex which brings our SFMS into agreement with ZFOURGE and MOSDEF. These systematic differences in SFRs due to using different conversion relations and dust laws highlights the need to identify a more robust method of measuring SFRs at $z > 1$ (*e.g.* Reddy et al. 2015; Shivaee et al. 2016).

In our analysis, we also compare star-forming galaxies that lie above, on, and below the star-forming main sequence (hereafter SFMS) as measured by $H\alpha$ emission. Using the best-fit to the combined cluster and field sample (Eq. 7), we calculate a galaxy's offset from the $H\alpha$ SFMS given its stellar mass. Because the typical scatter in the $H\alpha$ SFMS is ~ 0.2 dex, we use $\Delta\text{SFR}(M_*) = 0.2$ dex to separate galaxies into those above (20), on (45), and below (18) the SFMS. Galaxies in these three classes (+SFMS, =SFMS, -SFMS) span the full range in stellar mass (Fig. 6, right).

The LIRGs also span the full range in stellar mass and $H\alpha_{\text{star}}$ -SFR for both field and cluster galaxies, and the most massive galaxies [$\log(M_*/M_\odot) \gtrsim 10$] tend to be LIRGs (Fig. 6, left). The LIRGs at $z \sim 2$ follow the same trend of increasing $H\alpha_{\text{star}}$ -SFR with stellar mass (Fig. 6; slope ~ 0.80), a somewhat surprising result given the large scatter

when comparing SFRs derived from $H\alpha_{\text{star}}$ to L_{IR} (see §3.2). LIRGs lie above, on, and below the star-forming main sequence as defined by their $H\alpha_{\text{star}}$ -SFRs (Fig. 6, right).

3.4. Galaxy Size-Stellar Mass Relation

How galaxy size correlates with stellar mass depends on galaxy type, *e.g.* quiescent galaxies with Sérsic indices of $n \sim 4$ tend to be smaller at a given stellar mass compared star-forming galaxies with $n \sim 1$ (Shen et al. 2003). With a limited spectroscopic sample of galaxies, Law et al. (2012) showed that the galaxy size-mass relation evolves with redshift. Most recently, van der Wel et al. (2014) used high resolution imaging from *Hubble Space Telescope* and photometric redshifts for $\sim 31,000$ galaxies to measure how the $r_{\text{eff}}-M_{\star}$ relation of star-forming galaxies has evolved since $z \sim 3$.

We measure Sérsic indices and effective radii for 83 of the 90 galaxies in our sample (see §2.3 and Table 1). We find that our $H\alpha$ -emitting $z \sim 2$ galaxies follow the same trend of increasing galaxy size with stellar mass measured by van der Wel et al. (2014) for galaxies at this epoch (Fig. 4). Most of our fitted galaxies (71 of 83) have Sérsic indices of $n \leq 2$, and most (80 of 83) have effective radii of $0.7 < r_{\text{eff}} < 5$ kpc (Fig. 7).

3.4.1. Cluster vs. Field

We find no difference in the galaxy size-stellar mass relation with environment for $H\alpha$ -emitting galaxies. The cluster and field populations have the same size distributions with similar average effective radii of $r_{\text{eff}} \sim 2.5 \pm 0.2$ kpc and $r_{\text{eff}} \sim 2.2 \pm 0.2$ kpc, respectively (Fig. 7). Least-squares fits to the $r_{\text{eff}}-M_{\star}$ distribution for the cluster and field populations confirms that, within the errors, the least-squares fits agree with the van der Wel et al. (2014) size-mass relation.

The astute reader may notice possible conflict with our results in Allen et al. (2015) reporting that star-forming cluster galaxies are $\sim 12\%$ larger than in the field. However, we do find evidence that at *fixed stellar mass*, our cluster galaxies are ~ 0.1 dex larger which is consistent with Allen et al. (2015). We refer to §3.4.4 below for details.

3.4.2. IR-Luminous Galaxies

IR-luminous galaxies (LIRGs) have different physical size and stellar mass distributions relative to the low-IR population. A K-S test of the size distributions (Fig. 7) confirms with $> 3\sigma$ significance that the LIRGs are larger with a median $r_{\text{eff}} \sim 3.8$ kpc compared to ~ 2.0 kpc for the low-IR galaxies (typical errors for both are ~ 0.3 kpc). LIRGs also are ~ 5 times more massive with $\log(M_{\star}/M_{\odot}) \sim 10.4$ compared to ~ 9.6 for the low-IR galaxies (Figs. 4 & 6). Even if we consider only galaxies with $\log(M_{\star}/M_{\odot}) > 9.6$, LIRGs and low-IR galaxies have statistically different absolute r_{eff} distributions.

The size difference between our LIRGs and the low-IR galaxies at $z \sim 2$ seem to be in conflict with Swinbank et al. (2010) who, using *Hubble Space Telescope*/WFC3/F160W imaging of 25 submm galaxies at $\bar{z} \sim 2.1$, find their submm galaxies have the same sizes as field galaxies at $1 < z < 3.5$ (both have typical half-light radii of $\sim 2.5-2.8$ kpc). We find our LIRGs are typically $\sim 70\%$ larger than the low-IR population (see also Kartaltepe et al. 2012). This discrepancy is likely due to our IR comparison being based on a mass-selected sample that identifies LIRGs to $\log(M_{\star}/M_{\odot}) \sim 9.5$ (Fig. 6) while Swinbank et al. (2010) is limited to galaxies with $\log(M_{\star}/M_{\odot}) > 10$, *i.e.* galaxies that are large regardless of their L_{IR} emission because they are massive.

3.4.3. Above, On, & Below the $H\alpha$ Star-Forming Main Sequence

Galaxies above, on, and below the $H\alpha$ star-forming main sequence (SFMS; see Fig. 6, right) also follow the same general trend of increasing galaxy size with stellar mass (Fig. 4, right). K-S tests confirm that the size distributions for all three groups are likely drawn from the same parent population.

One concern in using $H\alpha$ SFRs obtained with slit spectroscopy is that we are biased towards compact star-forming galaxies, *e.g.* significant slit losses in the spectroscopic flux measurements will cause smaller galaxies to appear to have higher $H\alpha$ -SFRs compared to larger galaxies. However, the slit-width of $0.7''$ is comparable to the typical effective radius of most of the galaxies ($r_{\text{eff}} \sim 0.35''$; Fig. 4). Most importantly, we flux calibrate our spectroscopic measurements using total galaxy fluxes anchored in deep ground and space-based photometry and confirm that the uncertainty in the spectrophotometric calibration is 0.08 mag (see §2.7 in Nanayakkara et al. 2016).

3.4.4. Galaxy Size at Fixed Stellar Mass

To identify more subtle differences in galaxy size at fixed stellar mass, we first fit a (2σ clipped) least squares to $r_{\text{eff}}-M_{\star}$ using our combined cluster and field sample:




Figure 7. *Top:* $H\alpha$ -selected cluster and field galaxies at $z \sim 2$ have the same size distribution as measured by the effective radius (r_{eff}); medians are shown as vertical lines. *Middle:* However, in the combined sample, the IR-luminous galaxies ($L_{\text{IR}} > 2 \times 10^{11} L_{\odot}$) tend to be ~ 0.25 dex larger ($\sim 70\%$ larger in linear space) than the low-IR galaxies. A K-S test confirms at $> 3\sigma$ significance that the LIRGs and low-IR galaxies have different size distributions. The LIRGs also tend to be more massive (see Fig. 6). *Bottom:* Galaxies above, on, and below the $H\alpha$ star-forming main sequence span a similar range in galaxy size, but +SFMS galaxies tend to have smaller r_{eff} at a given stellar mass compared to -SFMS galaxies (Fig. 4).

$$\Delta[\log(r_{\text{eff}}, M_{\star})] = \log(r_{\text{eff}}, M_{\star}) - [(0.253 \times M_{\star}) - 2.12] \quad (8)$$

Our fitted least squares is virtually the same as the relation measured by [van der Wel et al. \(2014\)](#) for galaxies at $z = 2.0$ (Fig. 4, right).

When controlling for stellar mass, we find that the $\Delta[\log(r_{\text{eff}}, M_{\star})]$ distributions for the cluster and field galaxies are likely drawn from different parent populations (Fig. 8, top; $p = 0.01$); this is in contrast to no difference in their *absolute* r_{eff} distributions (Fig. 7). At fixed $M_{\star}$, $H\alpha$ -emitting cluster galaxies are ~ 0.1 dex larger than their field counterparts. Our result is consistent with [Allen et al. \(2015\)](#) who find that star-forming cluster galaxies as identified by their UVJ colors are $\sim 12\%$ larger than those in the field.

There is also a higher likelihood that at fixed stellar mass, galaxies above the SFMS are drawn from a different $\Delta[\log(r_{\text{eff}}, M_{\star})]$ parent population than those below (Fig. 8, bottom; $p = 0.05$). The +SFMS galaxies are ~ 0.1 dex smaller at a fixed $M_{\star}$ compared to -SFMS galaxies (Fig. 4). The compact nature of the +SFMS galaxies across the




Figure 8. The same as Fig. 7 but showing the difference in r_{eff} at a fixed stellar mass. Here $\Delta[\log(r_{\text{eff}}, M_*)]$ is determined using the (2σ clipped) least squares fit to $r_{\text{eff}}-M_*$ of our combined cluster and field galaxies (Fig. 4, black line in right panel). Compared to their absolute r_{eff} distributions (Fig. 7), K-S tests now measure higher likelihoods that the cluster and field galaxies are drawn from different $\Delta[\log(r_{\text{eff}}, M_*)]$ parent populations ($p = 0.01$); this is also true for the above vs. below SFMS galaxies ($p = 0.05$). The $\Delta[\log(r_{\text{eff}}, M_*)]$ distributions of the low-IR and LIRGs are more similar ($p = 0.06$).

entire stellar mass range suggests that their star formation is more centralized than in the $-$ SFMS galaxies (see also §4.2).

A K-S test of the $\Delta[\log(r_{\text{eff}}, M_*)]$ distributions for the low-LIR vs. LIRGs measures $p = 0.06$ which is not as statistically significant as when comparing their absolute r_{eff} distributions ($p = 9.6 \times 10^{-6}$). Because LIRGs are more massive (Fig. 6), they also tend to have larger radii. Thus controlling for stellar mass reduces differences in the LIRG and low-LIR populations.

3.5. Galaxy Morphology & Stellar Ages

Having measured Sérsic indices for 83 galaxies in our $\text{H}\alpha$ -emitting sample, we can compare the galaxy morphologies of the different populations. We find that all the galaxy populations (field vs. cluster, LIRG vs. low-IR, above/on/below SFMS) have comparable distributions in Sérsic index as measured by a K-S test. Most of the galaxies (71/83) are disk-dominated systems ($n \leq 2$).

The SED-based ages from ZFOURGE (Straatman et al. 2016) confirm that the cluster and field galaxies have similar age distributions of ~ 8.5 Gyr. This is also true for the LIRG and low-IR populations (both are ~ 8.5 Gyr).




Figure 9. *Left:* There are no differences in the cluster (filled red circles) and field (filled blue stars) galaxies when comparing their $H\alpha_{\text{star}}$ star formation rate to their WFC3/F160W galaxy size. The solid line in both panels is the least-squares fit (2σ outliers removed) to the combined sample. LIRGs (open orange stars) tend to be larger than low-IR galaxies in both environments. *Right:* Galaxies above (filled squares), on (open crosses), and below (filled triangles) the $H\alpha$ SFMS (see Fig. 6, right) populate different regions: +SFMS galaxies have higher $H\alpha_{\text{star}}$ -SFRs at a given size compared to -SFMS galaxies.

However, comparison of the galaxies above (+SFMS), on (=SFMS), and below (-SFMS) galaxies shows that their average stellar ages increases from ~ 8.3 , ~ 8.6 , to ~ 8.7 Gyr respectively. The younger light-weighted stellar ages of the +SFMS galaxies is consistent with a starburst nature.

3.6. Spatial Extent of $H\alpha_{\text{star}}$ -Star Formation

Using the star formation rates derived from $H\alpha_{\text{star}}$, the effective radii measured using WFC3/F160W imaging, and stellar masses from SED fitting, we first compare the $H\alpha_{\text{star}}$ -SFR to galaxy size (r_{eff} , Fig. 9; see §2.3 and Table 1). Our assumption that the $H\alpha$ radii are comparable to the rest-frame optical radii is supported by results from Förster Schreiber et al. (SINS; 2011) who combined IFU and HST observations of six $H\alpha$ -emitting galaxies at $z \sim 2$ and find no significant differences in their sizes and structural parameters at these wavelengths.

The cluster and field galaxies have similar distributions, and least-squares fits (2σ outliers removed) confirm that both populations have the same slopes within the errors. As seen in Fig. 7, the LIRGs tend to have larger r_{eff} than the low-IR galaxies because the LIRGs are more massive. In contrast, galaxies above the SFMS have higher $H\alpha_{\text{star}}$ -SFRs at a given size compared to those below the SFMS (Fig. 9).

We find similar results when comparing the star formation rate surface density [$\Sigma(H\alpha_{\text{star}})$; see Eq. 4] to galaxy size (r_{eff} , Fig. 10) and stellar mass (M_* , Fig. 11). The cluster and field galaxies have similar distributions, and least-squares fits (2σ -clipped) to $\Sigma(H\alpha_{\text{star}})$ - r_{eff} and $\Sigma(H\alpha_{\text{star}})$ - M_* confirm that both populations have the same slopes within the errors. Note that our sample spans a range in galaxy size [$0.5 < r_{\text{eff}}$ (kpc) < 8], star formation rate surface density [$0.01 < \Sigma(H\alpha_{\text{star}}) < 5$] where the units are $M_\odot \text{ yr}^{-1} \text{ kpc}^{-2}$, and stellar mass [$9 < \log(M_*/M_\odot) < 11$].

In contrast, the LIRGs and low-IR populations are different: at a given galaxy size, LIRGs tend to have higher SFR surface densities (Fig. 10, left). As noted in §3.3.2, LIRGs also are typically ~ 5 times more massive (Fig. 11) and physically larger by $\sim 70\%$. However, LIRGs are not all starbursts, *i.e.* LIRGs are found above, on, and below the SFMS (Fig. 6).

If we consider instead galaxies that lie above the SFMS, these +SFMS systems have higher SFR surface densities than -SFMS galaxies (Fig. 10, right). At a given stellar mass, the +SFMS galaxies tend to have smaller radii (Fig. 4) and higher $\Sigma(H\alpha_{\text{star}})$ (Fig. 11) compared to galaxies on/below the SFMS. Our results suggest that the $H\alpha$ star formation in +SFMS is more concentrated than those on/below the SFMS.

3.7. RHAPSODY-G Simulations: SFR- M_*




Figure 10. *Left:* The star formation rate surface density $\Sigma(\text{H}\alpha_{\text{star}})$ is measured with $\text{H}\alpha_{\text{star}}$ star formation rate and WFC3/F160W galaxy size, and the solid line is the least-squares fit (2σ outliers removed) to the combined sample. Cluster (filled circles) and field (line stars) galaxies have the same distribution in $\Sigma(\text{H}\alpha_{\text{star}})$ - r_{eff} . In contrast, the LIRGs (open stars) tend to be larger and have higher $\Sigma(\text{H}\alpha_{\text{star}})$ compared to low-IR galaxies, *i.e.* massive star-forming galaxies tend to have larger r_{eff} and also be LIRGs. *Right:* Galaxies above (filled squares), on (open crosses), and below (filled triangles) the $\text{H}\alpha$ SFMS (see Fig. 6, right) populate different regions: +SFMS galaxies are forming stars more intensely than -SFMS galaxies across the range in galaxy size.




Figure 11. *Left:* The star formation rate surface density $\Sigma(\text{H}\alpha_{\text{star}})$ compared to stellar mass M_* where the solid line is the least-squares fit (2σ outliers removed) to the combined sample. Cluster (filled circles) and field (line stars) galaxies have the same distribution in $\Sigma(\text{H}\alpha_{\text{star}})$ - M_* . LIRGs (open stars) are more massive than low-IR galaxies, but both populations span the range in $\Sigma(\text{H}\alpha_{\text{star}})$. *Right:* Galaxies above (filled squares), on (open crosses), and below (filled triangles) the $\text{H}\alpha$ SFMS (see Fig. 6, right) populate different regions: +SFMS galaxies are forming stars more intensely than -SFMS galaxies across the range in stellar mass.

We compare our measured $H\alpha_{\text{star}}$ star formation rate versus stellar mass relation to predictions from the RHAPSODY-G simulations of massive galaxy clusters ($M_{\text{vir}} > 6 \times 10^{14} M_{\odot}$ at $z = 0$; [Hahn et al. 2015](#); [Martizzi et al. 2016](#)). These cosmological hydrodynamical zoom-in simulations (R4K resolution) use the RAMSES adaptive mesh refined (AMR) code ([Teyssier 2002](#)) to reach a spatial resolution of $3.8h^{-1}$ kpc (physical), dark matter particle mass resolution of $8.22 \times 10^8 h^{-1} M_{\odot}$, and baryonic mass resolution of $1.8 \times 10^8 h^{-1} M_{\odot}$. The simulations assume the standard Λ CDM cosmology ($\Omega_M = 0.25$, $\Omega_{\Lambda} = 0.75$, $\Omega_b = 0.045$, $h = 0.7$) and include gas cooling, star formation, metal enrichment, and feedback from supernovae and AGN.

The RHAPSODY-G cluster simulations are well-matched to our COSMOS cluster at $z_{\text{cl}} = 2.1$. As detailed in [Yuan et al. \(2014, see their §4\)](#), its measured velocity dispersion of $\sigma_{1D} = 552 \text{ km s}^{-1}$ corresponds to a virial mass of $\log(M_{\text{vir}}/M_{\star}) \sim 13.5$. Merger trees from the GigggleZ Gpc simulation ([Poole et al. 2015](#)) shows that such systems grow into a Virgo-mass cluster with $\log(M_{\text{vir}}/M_{\star}) \sim 14.4$ by $z \sim 0$.

We consider only simulated cluster galaxies at $z = 2$ with star formation rates $> 1 M_{\odot} \text{ yr}^{-1}$; these galaxies have stellar masses of $\log(M_{\star}/M_{\odot}) = 9 - 12$. Here we assume that selecting by SFR is equivalent to the instantaneous observed SFR as measured by $H\alpha_{\text{star}}$. We cannot apply the same observed UVJ selection as rest-frame colors are not available for the simulated galaxies.

From three RHAPSODY-G cluster realizations, the least-squares fit to the SFR- $M_{\star}$ relation is:

$$\log[\text{SFR}(M_{\odot} \text{ yr}^{-1})] = 1.08 \{\log[M_{\star}(M_{\odot})] - 10\} + \log(4.5) \quad (9)$$

The RHAPSODY-G slope to the SFR- $M_{\star}$ relation is steeper than that of the observed galaxies at $z \sim 2$: 1.08 vs. 0.61 (Fig. 6, right panel: gray and cyan lines respectively). Although the slopes are consistent within the scatter of the simulations and observations (see §3.3), the SFRs predicted by RHAPSODY-G are lower by a factor of ~ 2 for most of the observed galaxies. This difference between predicted and observed SFRs at a given stellar mass (*i.e.* the specific SFR) is known to exist for field comparisons (*e.g.* [Davé et al. 2016](#)). Here we show that this discrepancy extends to the cluster environment as well, *i.e.* simulations over-predict how efficiently galaxies quench at a given stellar mass for both the cluster and field environments (see [Somerville & Davé 2015](#), and references therein). In the case of RHAPSODY-G, star formation histories at high redshift are slightly under-resolved due to the mass resolution. Future simulations with higher resolution combined with multi-epoch observations are needed to improve galaxy formation modeling at $z \sim 2$. We will explore more key scaling relations compared to simulations in future work.

4. DISCUSSION

Our analysis focuses on comparing at $z \sim 2$ (i) cluster to field galaxies (37 vs. 53); (ii) galaxies with $L_{\text{IR}} > 2 \times 10^{11} L_{\odot}$ (LIRGs) to the low-IR population (21 vs. 69); and (iii) galaxies above, on, and below the $H\alpha$ star-forming main sequence (SFMS). Because ours is a stellar mass-selected sample of $H\alpha$ -emitting galaxies [$\log(M_{\star}/M_{\odot}) > 9$; see Fig. 6], we are not limited to the high mass end of the galaxy population. We consider only $H\alpha$ -selected galaxies at $1.9 < z < 2.4$ because the redshifts for the quiescent galaxies are based on photometry and/or grism spectroscopy ([Tomczak et al. 2014](#); [Momcheva et al. 2015](#)), neither of which are as precise as our Keck/MOSFIRE redshifts determined with $H\alpha$. We confirm that selecting field galaxies using a more stringent cut of $> 8\sigma_{1D}$ from the mean cluster redshift does not change our results.

4.1. $H\alpha$ -Emitting Galaxies: Little Evidence of Environmental Dependence at $z \sim 2$

Our original motivation was to quantify how galaxy properties vary with environment at $z \sim 2$. However, we find little evidence for environmental dependence in $H\alpha$ -emitting galaxies at $z \sim 2$. We consistently measure the same relations for cluster and field galaxies when comparing their $H\alpha_{\text{star}}$ star formation rate to stellar mass (Fig. 6), galaxy size to stellar mass (Figs. 4 & 7), and star formation concentration (Fig. 9, 10, & 11). The fraction of LIRGs and their median L_{IR} are also the same in the cluster and field (§3.1). In our study, the only measureable difference is that at fixed stellar mass, $H\alpha$ -emitting cluster galaxies are ~ 0.1 dex larger than the field (Fig. 8).

In terms of their physical properties, the $H\alpha$ -emitting cluster galaxies at $z_{\text{cl}} = 2.1$ are essentially the same population as the field. This is consistent with our results in [Kacprzak et al. \(2015\)](#) showing that these very same cluster and field galaxies also follow the same gas-phase metallicity versus stellar mass relation (MZR). In addition, we find no evidence for an environmental dependence when comparing their kinematic scaling relations ([Alcorn et al. 2016](#); [Straatman et al. 2016](#)).

The handful of existing studies on galaxy overdensities at $z \gtrsim 2$ similarly find little evidence for environmental effects. Using narrow-band imaging, [Koyama et al. \(2013\)](#) measure the same SFR- $M_{\star}$ relation for $H\alpha$ -emitters in a

$z = 2.16$ proto-cluster as in the field. Using high resolution imaging from the *Hubble Space Telescope*, [Peter et al. \(2007\)](#) measure the same size (radius) distributions for field and proto-cluster galaxies at $z = 2.3$.

In contrast, [Papovich et al. \(2012\)](#) find that quiescent cluster galaxies at $z = 1.62$ are larger than their field counterparts, and [Quadri et al. \(2012\)](#) find a higher fraction of quiescent galaxies in the same cluster. Several studies also find evidence of enhanced star formation in cluster galaxies at $z < 2$ ([Tran et al. 2010](#); [Brodwin et al. 2013](#); [Santos et al. 2014](#); [Tran et al. 2015](#); [Webb et al. 2015](#)). The lack of convincing evidence for strong environmental effects at $z \gtrsim 2$ combined with the increasing differences between cluster and field galaxies at lower redshifts pinpoints to $1.5 \lesssim z \lesssim 2$ as the critical epoch for ending star formation in cluster galaxies and building the spheroid population in clusters.

4.2. Tracking Galaxy Growth with the $H\alpha$ Star-Forming Main Sequence

Given the physical properties of $H\alpha$ -emitting galaxies show little environmental dependence (see above), we can use the combined cluster and field sample at $z \sim 2$ to compare galaxies above, on, and below the star-forming main sequence (SFMS) as well as compare the IR-luminous (21; LIRG) to low-IR (69) populations (Figs. 6 through 11). Because our spectroscopic target selection is based on ZFOURGE, we are mass-limited to $\log(M_*/M_\odot) \sim 9$ at $z \sim 2$ ([Tomczak et al. 2014](#); [Nanayakkara et al. 2016](#)).

Being a LIRG does not necessarily mean the galaxy is a starburst because LIRGs are found above, on, and below the SFMS (Fig. 6, left). Rather, IR-luminosity tends to track closely with stellar mass such that massive galaxies [$\log(M_*/M_\odot) > 10$] tend to be LIRGs. On average, LIRGs are ~ 5 times more massive and $\sim 70\%$ larger than low-IR galaxies (Figs. 4, 7, & 9). When controlling for stellar mass, there is less difference in the size distributions the LIRGs and low-IR galaxies (Fig. 8). Note that the mass range of our $H\alpha$ -emitting galaxies reaches $\log(M_*/M_\odot) \sim 9$, *i.e.* a factor of about 5 – 10 times lower than previous studies that compared LIRGs to the general galaxy population at $z > 1$ (*e.g.* [Swinbank et al. 2010](#)).

In terms of tracking how galaxies grow, systems that lie above the $H\alpha$ star-forming main sequence (+SFMS) have smaller radii at a given stellar mass than those that are below the SFMS (Fig. 4, right; see §3.4.3). The +SFMS galaxies tend to have higher $H\alpha_{\text{star}}$ -SFRs at a given galaxy size (Fig. 9) and higher $H\alpha_{\text{star}}$ SFR surface densities compared to those below [$\Sigma(H\alpha_{\text{star}})$; Figs. 10 & 11], *i.e.* their star formation is more compact. The +SFMS galaxies also have younger SED-based stellar ages of ~ 8.3 Gyr compared to ~ 8.7 Gyr for –SFMS galaxies. Taken as a whole, our results indicate that +SFMS are starbursts with $H\alpha$ star formation concentrated in their cores (see also [Barro et al. 2015](#)).

At $z \sim 1$, field galaxies are preferentially growing their disks ([Nelson et al. 2016](#)). In combination with our observations indicating that starbursts at $z \sim 2$ are growing their stellar cores, these results suggest a sequence where +SFMS galaxies are building up their stellar cores at $z \sim 2$ and then their stellar disks at $z \sim 1$, *i.e.* inside-out growth, likely by continuing gas accretion at $z < 2$ (*e.g.* [Kacprzak et al. 2016](#)). Such a scenario naturally produces the older stellar populations of bulges relative to disk. This can also explain the rise of spheroids in clusters if the cluster environment prevents the growth of stellar disks even as star formation in the galaxies' cores is quenched at $z < 1.5$ ([Brodwin et al. 2013](#); [Tran et al. 2015](#)). While our hypothesis is based on the +SFMS galaxies, we note that galaxies at $z \sim 2$ in general must grow physically larger by $z \sim 1$ (*e.g.* [van der Wel et al. 2014](#)).

4.3. Star Formation Rates at $z \sim 2$: Caveat Emptor

Our analysis is based on the relative comparison of cluster and field galaxies where properties for both are determined in the same manner. Thus our results do not depend on the absolute conversion of, *e.g.* $H\alpha$ flux to SFR. However, we do find that the $H\alpha_{\text{star}}$ SFRs are offset from L_{IR} SFRs (Fig. 5). The large uncertainty and likely offset from relations measured at $z \sim 0$ brings into question our ability to measure reliable SFRs at $z > 1$.

There are several ongoing efforts to better understand star formation and dust laws at $z > 1$ that should help with calibrating existing relations. Recent studies at $z \sim 2$ find evidence of changing ionization conditions ([Sanders et al. 2016](#)) as well as different dust laws ([Reddy et al. 2015](#); [Forrest et al. 2016](#); [Shivaei et al. 2016](#)) that can be incorporated into models. However, until we identify a more robust method for measuring SFRs in the distant universe, direct comparisons between studies will require carefully accounting for different methods of measuring SFRs.

5. CONCLUSIONS

Our ZFIRE program combines Keck/MOSFIRE spectroscopy with the wealth of multi-wavelength observations available in the COSMOS legacy field to explore galaxy scaling relations as a function of environment at $z \sim 2$. Our advantage is that we select galaxies at $z \sim 2$ based on their stellar masses as measured by ZFOURGE, a deep imaging

survey that uses medium-band NIR filters to obtain high-precision photometric redshifts ($\sigma_z \sim 0.02$; [Straatman et al. 2016](#)). We focus on the spectroscopically confirmed galaxy cluster at $z = 2.095$ in the COSMOS legacy field ([Spitler et al. 2012](#); [Yuan et al. 2014](#)) and compare to the field population at $z \sim 2$.

In comparing H α -emitting cluster (37) and field (53) galaxies [$\log(M_*/M_\odot) > 9$; AGN removed], we find little evidence of environmental influence on any of the galaxy scaling relations. Both cluster and field populations are consistent with published relations between star formation rate and stellar mass (SFR- M_* ; Fig. 6) as well as galaxy size and stellar mass ($r_{\text{eff}}-M_*$; Fig. 4) at $z \sim 2$. The cluster and field populations also have the same distribution when comparing their H α_{star} -SFR surface density [$\Sigma(\text{H}\alpha_{\text{star}})$] to galaxy size and stellar mass (Figs. 10 & 11). The results in this analysis mirror our existing ZFIRE results that show these same cluster and field galaxies have the same gas-phase metallicity vs. stellar mass relation ([Kacprzak et al. 2015](#)), kinematic mass vs. stellar mass ([Alcorn et al. 2016](#)), and ISM conditions ([Kewley et al. 2016](#)). The only subtle indication of possible environmental dependence is that at fixed stellar mass, the H α -emitting cluster galaxies are ~ 0.1 dex larger than in the field (Fig. 8).

Using *Spitzer*/24 μm observations, we identify 21 galaxies with $L_{\text{IR}} > 2 \times 10^{11} L_\odot$, *i.e.* Luminous Infra-Red Galaxies (LIRGs). Note that our mass range of $\log(M_*/M_\odot) \sim 9$ is a factor of about 5 – 10 times lower than previous studies that compared LIRGs to the general galaxy population at $z > 1$ (*e.g.* [Swinbank et al. 2010](#)). The LIRG fraction is comparable within errors between the cluster and the field (19% and 26% respectively), and we do not find any indication that LIRGs in the cluster are different from those in the field. IR-luminosity tracks with stellar mass such that our most massive galaxies [$\log(M_*/M_\odot) > 10$] are dominated by LIRGs. As a result, LIRGs tend to be ~ 5 times more massive with radii that are $\sim 70\%$ larger than low-IR galaxies ($r_{\text{eff}} \sim 3.8$ kpc vs. ~ 2.0 kpc; Fig. 7). The LIRGs are not all starbursts as they are found above, on, and below the H α Star-Forming Main Sequence (SFMS; Fig. 6).

We show that separating galaxies into those above the H α Star-Forming Main Sequence (+SFMS), on, and below (–SFMS) provides insight into how galaxies grow (Fig. 6). Galaxies in the three groups span the full range in parameter space, but the +SFMS galaxies have smaller radii at a given stellar mass compared to –SFMS (Fig. 4). The +SFMS galaxies also tend to have higher SFR surface densities compared to galaxies with depressed SFRs (Figs. 9, 10, & 11), and younger SED-based stellar ages compared to galaxies below the SFMS (~ 8.3 Gyr vs. ~ 8.7 Gyr).

These lines of evidence indicate that +SFMS galaxies (starbursts) have concentrated H α star formation and are actively growing their cores at $z \sim 2$. We infer that while starbursts in the field go on to grow their stellar disks at $z \sim 1$ ([Nelson et al. 2016](#)), cluster starbursts are likely to be quenching their star formation at $z < 2$ ([Brodwin et al. 2013](#); [Tran et al. 2015](#)) to then evolve into quiescent spheroids ([Quadri et al. 2012](#); [Papovich et al. 2012](#)).

We compare the H α SFR- M_* relation to predictions from the RHAPSODY-G simulations of massive galaxy clusters ($> 6 \times 10^{14} M_\odot$ at $z = 0$) based on the RAMSES Adaptive Mesh Refinement code. We find that the predicted slope for the SFR- M_* relation is steeper than the observed values (1.08 vs. 0.61), and that the predicted SFRs are ~ 2 times lower than observed. Simulations in general continue to over-predict how efficiently galaxies quench at a given stellar mass in both the cluster and field environments. We will continue to explore how observed galaxy scaling relations compare to simulations in future work.

On a cautionary note, there is considerable scatter and likely offset in star formation rates based on H α and those based on IR luminosity (or UV+IR) at $z \sim 2$ (Fig. 5). This is in contrast to the relatively small scatter at $z \sim 0$ between H α and 24 μm derived star formation rates for activity at $< 100 M_\odot \text{ yr}^{-1}$ (*e.g.* [Hao et al. 2011](#)). It is sobering to consider the large uncertainty in measuring robust SFRs, especially at higher redshifts when SFRs are increasing in general ([Garn & Best 2010](#); [Whitaker et al. 2014](#); [Tomczak et al. 2016](#)). Our ability to accurately measure star formation at $z \gtrsim 2$ is likely to be limited due to, *e.g.* our understanding of how ionization conditions evolve. However, we stress that the strength of this analysis lies in using the same observables to directly compare across different galaxy populations at $z \sim 2$.

In a companion ZFIRE paper, we estimate gas masses and gas depletion timescales for the same cluster and field galaxies at $z \sim 2$. Ongoing analyses also include a comparison of the Tully-Fisher relation ([Straatman et al.](#), submitted) and constraints on the Initial Mass Function ([Nanayakkara et al.](#), in prep). By measuring galaxy scaling relations for cluster and field galaxies at $z \sim 2$, ZFIRE provides a unique benchmark for quantifying galaxy evolution as a function of environment.

Table 1. Galaxy Properties^a

ZFIRE ^b	ZFOURGE ^b	$\alpha(2000)$	$\delta(2000)$	z_{spec}	fH α ^c	err(fH α) ^c	log(L _{IR} /L $\odot$) ^d	log(M $_{\star}$ /M $\odot$)	AV,star	log(t_{star}) ^e	SFR(H α_{star}) ^f	Sérsic n	r _{eff} (")	Pflag ^g
237	912	150.19057	2.18848	2.1572	1.46	0.17	...	9.65	0.6	8.1	6.0	...	...	-99
342	1108	150.19051	2.19065	2.1549	3.98	0.09	11.93	10.45	1.1	8.9	31.3	0.8	0.4	0
1085	2114	150.18338	2.20192	2.1882	2.53	0.07	...	9.60	0.1	8.4	5.6	1.3	0.2	0
1180	2168	150.12984	2.20287	2.0976	1.33	0.15	...	8.94	0.0	8.1	2.3	4.0	0.2	2
1349	2517	150.20306	2.20554	2.1888	1.04	0.06	11.23	9.82	0.3	8.3	3.0	4.0	0.1	1
1385	2510	150.12344	2.20565	2.0978	3.28	0.23	...	9.30	0.1	8.0	6.5	0.5	0.3	0
1617	2989	150.09697	2.20917	2.1732	1.32	0.14	11.22	10.14	0.5	8.7	4.8	0.9	0.7	2
1814	3175	150.16809	2.21129	2.1704	4.55	0.10	...	9.95	0.4	8.5	14.6	1.0	0.3	0
2007	3375	150.16566	2.21366	2.0086	1.17	0.14	11.17	9.42	0.4	8.5	3.1	0.9	0.1	0
2153	3669	150.16533	2.21584	2.0123	4.86	0.18	11.48	10.07	0.7	8.8	19.2	4.0	0.9	2
2522	4084	150.19379	2.22011	2.1511	1.10	0.11	11.48	9.64	0.3	8.1	3.0	0.5	0.4	0

^a The complete version of this table is available in electronic format.

^b We list galaxy identification numbers from ZFIRE (Nanayakkara et al. 2016) and ZFOURGE (Straatman et al. 2016). We include only galaxies with spectroscopic redshift quality flag of $Q_z = 3$ (Nanayakkara et al. 2016) and $1.9 < z_{\text{spec}} < 2.4$. Cluster members have $2.08 < z_{\text{spec}} < 2.12$ (Yuan et al. 2014).

^c Observed H α fluxes and errors are in units of 10^{-17} erg s $^{-1}$ cm $^{-2}$.

^d In our analysis of IR-luminous vs. low-IR systems, we select IR-luminous galaxies using $\log(L_{\text{IR}}/L_{\odot}) > 11.3$.

^e Stellar age in units of Gyr and based on SED fitting with FAST (Kriek et al. 2009).

^f H α_{star} star formation rates in units of M $\odot$ yr $^{-1}$ and based on dust-corrected H α fluxes (Eq. 2; see §2.4).

^g Pflag denotes quality of profile fit used to measure the Sérsic index n and the effective radius r_{eff}. Pflag values are -99 (not fit), 0 (good fit), 1 (fair fit), and 2 (questionable fit).

We are grateful to the MOSFIRE team with special thanks to M. Kassis, J. Lyke, G. Wirth, and L. Rizzi on the Keck support staff. K. Tran thanks M. Kriek, A. Shapley, S. Price, B. Forrest, and Jimmy for helpful discussions. We also thank the referee for a thoughtful and constructive report. This work was supported by a NASA Keck PI Data Award administered by the NASA Exoplanet Science Institute. Data presented herein were obtained at the W. M. Keck Observatory from telescope time allocated to NASA through the agency's scientific partnership with the California Institute of Technology and the University of California. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation. K. Tran acknowledges that this material is based upon work supported by the National Science Foundation under Grant Number 1410728. GGK acknowledges the support of the Australian Research Council through the award of a Future Fellowship (FT140100933). The authors wish to recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.

REFERENCES

- Alcorn, L. Y., Tran, K.-V. H., Kacprzak, G. G., et al. 2016, *ApJL*, 825, L2
- Allen, R. J., Kacprzak, G. G., Spitler, L. R., et al. 2015, *ApJ*, 806, 3
- Barro, G., Faber, S. M., Koo, D. C., et al. 2015, *ArXiv e-prints*, arXiv:1509.00469
- Bassett, R., Papovich, C., Lotz, J. M., et al. 2013, *ApJ*, 770, 58
- Brammer, G. B., van Dokkum, P. G., & Coppi, P. 2008, *ApJ*, 686, 1503
- Brammer, G. B., van Dokkum, P. G., Franx, M., et al. 2012, *ApJS*, 200, 13
- Brodwin, M., Stanford, S. A., Gonzalez, A. H., et al. 2013, *ApJ*, 779, 138
- Brown, M. J. I., Moustakas, J., Smith, J.-D. T., et al. 2014, *ApJS*, 212, 18
- Calzetti, D., Armus, L., Bohlin, R. C., et al. 2000, *ApJ*, 533, 682
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Casey, C. M., Narayanan, D., & Cooray, A. 2014, *PhR*, 541, 45
- Chabrier, G. 2003, *PASP*, 115, 763
- Cowley, M. J., Spitler, L. R., Tran, K.-V. H., et al. 2016, *MNRAS*, 457, 629
- Davé, R., Oppenheimer, B. D., & Finlator, K. 2011, *MNRAS*, 415, 11
- Davé, R., Rafieferantsoa, M. H., Thompson, R. J., & Hopkins, P. F. 2016, *ArXiv e-prints*, arXiv:1610.01626
- Forrest, B., Tran, K.-V. H., Tomczak, A. R., et al. 2016, *ApJL*, 818, L26
- Förster Schreiber, N. M., Shapley, A. E., Erb, D. K., et al. 2011, *ApJ*, 731, 65
- Garn, T., & Best, P. N. 2010, *MNRAS*, 409, 421
- Genel, S., Vogelsberger, M., Springel, V., et al. 2014, *MNRAS*, 445, 175
- Genzel, R., Tacconi, L. J., Lutz, D., et al. 2015, *ApJ*, 800, 20
- Gobat, R., Strazzullo, V., Daddi, E., Onodera, M., & et al. 2013, *ApJ*, 776, 9
- Grogin, N. A., Kocevski, D. D., Faber, S. M., et al. 2011, *ApJS*, 197, 35
- Hahn, O., Martizzi, D., Wu, H.-Y., et al. 2015, *ArXiv e-prints*, arXiv:1509.04289
- Hao, C.-N., Kennicutt, R. C., Johnson, B. D., et al. 2011, *ApJ*, 741, 124
- Hung, C.-L., Casey, C. M., Chiang, Y.-K., et al. 2016, *ApJ*, 826, 130
- Ibar, E., Sobral, D., Best, P. N., et al. 2013, *MNRAS*, 434, 3218
- Kacprzak, G. G., van de Voort, F., Glazebrook, K., et al. 2016, *ApJL*, 826, L11
- Kacprzak, G. G., Yuan, T., Nanayakkara, T., et al. 2015, *ApJL*, 802, L26
- Kartaltepe, J. S., Dickinson, M., Alexander, D. M., et al. 2012, *ApJ*, 757, 23
- Kenney, J. D. P., & Koopmann, R. A. 1999, *AJ*, 117, 181
- Kennicutt, R. C. 1998, *ARA&A*, 36, 189
- Kennicutt, R. C., & Evans, N. J. 2012, *ARA&A*, 50, 531
- Kewley, L. J., Yuan, T., Nanayakkara, T., et al. 2016, *ApJ*, 819, 100
- Koopmann, R. A., & Kenney, J. D. P. 2004, *ApJ*, 613, 866
- Koyama, Y., Kodama, T., Tadaki, K.-i., et al. 2013, *MNRAS*, 428, 1551
- Kriek, M., van Dokkum, P. G., Labbé, I., et al. 2009, *ApJ*, 700, 221
- Kroupa, P. 2001, *MNRAS*, 322, 231
- Labbé, I., Bouwens, R., Illingworth, G. D., & Franx, M. 2006, *ApJL*, 649, L67
- Law, D. R., Steidel, C. C., Shapley, A. E., et al. 2012, *ApJ*, 745, 85
- Lutz, D. 2014, *ARA&A*, 52, 373
- Madau, P., & Dickinson, M. 2014, *ARA&A*, 52, 415
- Martizzi, D., Hahn, O., Wu, H.-Y., et al. 2016, *MNRAS*, 459, 4408
- McLean, I. S., Steidel, C. C., Epps, H. W., Konidaris, N., & et al. 2012, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 8446, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series
- Momcheva, I. G., Brammer, G. B., van Dokkum, P. G., et al. 2015, *ArXiv e-prints*, arXiv:1510.02106
- Muzzin, A., Wilson, G., Yee, H. K. C., Gilbank, D., & et al. 2012, *ApJ*, 746, 188
- Nanayakkara, T., Glazebrook, K., Kacprzak, G. G., et al. 2016, *ApJ*, 828, 21
- Nelson, E. J., van Dokkum, P. G., Förster Schreiber, N. M., et al. 2016, *ApJ*, 828, 27
- Noeske, K. G., Weiner, B. J., Faber, S. M., et al. 2007, *ApJL*, 660, L43
- Papovich, C., Bassett, R., Lotz, J. M., et al. 2012, *ApJ*, 750, 93
- Papovich, C., Momcheva, I., Willmer, C. N. A., et al. 2010, *ApJ*, 716, 1503
- Peng, Y.-j., Lilly, S. J., Kovač, K., et al. 2010, *ApJ*, 721, 193
- Persson, S. E., Murphy, D. C., Smee, S., et al. 2013, *PASP*, 125, 654
- Peter, A. H. G., Shapley, A. E., Law, D. R., et al. 2007, *ApJ*, 668, 23
- Poole, G. B., Blake, C., Marín, F. A., et al. 2015, *MNRAS*, 449, 1454

- Popesso, P., Biviano, A., Rodighiero, G., et al. 2012, *A&A*, 537, A58
- Price, S. H., Kriek, M., Brammer, G. B., Conroy, C., & et al. 2014, *ApJ*, 788, 86
- Quadri, R. F., Williams, R. J., Franx, M., & Hildebrandt, H. 2012, *ApJ*, 744, 88
- Reddy, N. A., Kriek, M., Shapley, A. E., et al. 2015, *ApJ*, 806, 259
- Rujopakarn, W., Rieke, G. H., Eisenstein, D. J., & Juneau, S. 2011, *ApJ*, 726, 93
- Saintonge, A., Tran, K.-V. H., & Holden, B. P. 2008, *ApJL*, 685, L113
- Sanders, D. B., & Mirabel, I. F. 1996, *ARA&A*, 34, 749
- Sanders, R. L., Shapley, A. E., Kriek, M., et al. 2015, *ApJ*, 799, 138
- . 2016, *ApJ*, 816, 23
- Santos, J. S., Altieri, B., Tanaka, M., et al. 2014, *MNRAS*, 438, 2565
- Schaye, J., Crain, R. A., Bower, R. G., et al. 2015, *MNRAS*, 446, 521
- Shen, S., Mo, H. J., White, S. D. M., et al. 2003, *MNRAS*, 343, 978
- Shivaei, I., Kriek, M., Reddy, N. A., Shapley, A. E., & et al. 2016, *ApJL*, 820, L23
- Skelton, R. E., Whitaker, K. E., Momcheva, I. G., et al. 2014, *ApJS*, 214, 24
- Somerville, R. S., & Davé, R. 2015, *ARA&A*, 53, 51
- Spitler, L. R., Labbé, I., Glazebrook, K., & et al. 2012, *ApJL*, 748, L21
- Steidel, C. C., Rudie, G. C., Strom, A. L., et al. 2014, *ApJ*, 795, 165
- Straatman, C. M. S., Spitler, L. R., Quadri, R. F., et al. 2016, *ApJ*, 830, 51
- Swinbank, A. M., Smail, I., Chapman, S. C., et al. 2010, *MNRAS*, 405, 234
- Teyssier, R. 2002, *A&A*, 385, 337
- Tomczak, A. R., Quadri, R. F., Tran, K.-V. H., et al. 2016, *ApJ*, 817, 118
- . 2014, *ApJ*, 783, 85
- Tonnesen, S., & Cen, R. 2014, *ApJ*, 788, 133
- Tran, K., Papovich, C., Saintonge, A., & et al. 2010, *ApJL*, 719, L126
- Tran, K.-V. H., Nanayakkara, T., Yuan, T., et al. 2015, *ApJ*, 811, 28
- van der Wel, A., Bell, E. F., Häussler, B., et al. 2012, *ApJS*, 203, 24
- van der Wel, A., Franx, M., van Dokkum, P. G., et al. 2014, *ApJ*, 788, 28
- Webb, T., Noble, A., DeGroot, A., et al. 2015, *ApJ*, 809, 173
- Wetzel, A. R., Tinker, J. L., & Conroy, C. 2012, *MNRAS*, 424, 232
- Whitaker, K. E., Franx, M., Bezanson, R., et al. 2015, *ApJL*, 811, L12
- Whitaker, K. E., Kriek, M., van Dokkum, P. G., & et al. 2012, *ApJ*, 745, 179
- Whitaker, K. E., Franx, M., Leja, J., et al. 2014, *ApJ*, 795, 104
- Williams, R. J., Quadri, R. F., Franx, M., van Dokkum, P., & Labbé, I. 2009, *ApJ*, 691, 1879
- Wuyts, S., Labbé, I., Förster Schreiber, N. M., et al. 2008, *ApJ*, 682, 985
- Wuyts, S., Labbé, I., Franx, M., et al. 2007, *ApJ*, 655, 51
- Wuyts, S., Förster Schreiber, N. M., van der Wel, A., et al. 2011, *ApJ*, 742, 96
- Yuan, T., Nanayakkara, T., Kacprzak, G. G., et al. 2014, *ApJL*, 795, L20
- Zeimann, G. R., Stanford, S. A., Brodwin, M., et al. 2012, *ApJ*, 756, 115